

Respite for 2-Dimensional Right Ventricular Imaging?

Rohan Samson, MD; Thierry H. Le Jemtel, MD

The complex geometry and structure of the right ventricular (RV) cavity make the assessment of RV function and volumes by 2-dimensional (2D) echocardiography arduous. Highly trabeculated endocardial border and bellows-like contractile pattern are not explicitly imaged by 2D echocardiography.^{1,2} Furthermore, the retrosternal position of the right ventricle hinders its visualization, and minimal changes in the angle of insonification affect RV chamber measurement and thereby, reproducibility.

See Article by Amsallem et al

Parameters of RV function and volume derived from 2D echocardiography (tricuspid annular plane systolic excursion, RV to left ventricular diameter ratio, and fractional area change) do not reliably predict the risk of RV failure during mechanical circulatory support with left ventricular assist device.^{3,4} Similarly, 2D assessment of RV function and volume in pulmonary hypertension has been challenging. The etiology and hemodynamic classification (pre- versus postcapillary) of pulmonary hypertension are important determinants of the RV remodeling process, leading to distinct RV remodeling phenotypes.^{5,6} In particular, the etiology of the pulmonary hypertension affects the RV deformation pattern and leads to contractile dysfunction involving the entire RV free wall or is localized to only the RV basal/apical regions.⁵

Predetermined geometric assumptions do not account for changes in RV geometry during the progression of the underlying condition. They jeopardize the accuracy of 2D quantification of RV function and volumes.⁷ Newer echocardiographic techniques that do not require geometric assumptions provide a far more reliable assessment of RV function and volumes than conventional 2D echocardiography.⁸ Three-dimensional echocardiography and speckle tracking imaging have been shown to predict RV failure and clinical outcome more accurately than 2D echocardiography in chronic pulmonary hypertension.^{5,7}

In this issue of *Circulation: Cardiovascular Imaging*, Amsallem et al⁹ propose a new 2D echocardiography-derived RV end-systolic remodeling index (RVESRI) in patients with

pulmonary arterial hypertension (PAH). Amsallem et al define the new RVESRI as the ratio of end-systolic RV lateral wall length to end-systolic septal height measured in the RV-focused apical 4-chamber view. The lateral wall length is measured from the lateral tricuspid annulus to the RV insertion of the interventricular septum. The septal height is measured as a straight line from the septal tricuspid annulus to the RV insertion on the interventricular septum. The RVESRI prognostic value was prospectively evaluated in 228 patients with PAH during a mean duration of 3.9±2.4 years in a single center. The primary end point of death, transplant, or hospitalization for heart failure was reached in 88 patients. The RVESRI was found to be more closely associated with clinical outcome than RV end-systolic size or the transverse to longitudinal RV ratio. When added to the Registry to Evaluate Early And Long-term PAH Disease Management score, the incremental predictive value of RVESRI was superior to that provided by 2D echocardiography-derived RV free wall longitudinal strain: C statistics of 0.81 (0.74 to 0.88) for Registry to Evaluate Early And Long-term PAH Disease Management score and 0.83 (0.77 to 0.89) for Registry to Evaluate Early And Long-term PAH Disease Management score+RVESRI. Thus, Amsallem et al advocate RVESRI as a simple reproducible prognostic marker in PAH.

Whether end-systolic RV free wall length to septal height ratio is a more accurate marker of RV remodeling in PAH than are conventional 2D echocardiographic parameters awaits further studies. As with other 2D echocardiographic measurements, an RV-focused apical 4-chamber view that avoids foreshortening, displays the largest basal RV diameter, and includes the entire RV free wall is essential for accurate quantification of RV linear measurements.¹⁰ Measurement of end-systolic RV free wall length relates to the longitudinal contraction of the right ventricle.¹¹ However, RV contractility does not only depend on the traction of the tricuspid annulus toward the apex. Bulging of the interventricular septum into the RV and radial displacement of the lateral free wall are important contributors to RV pump function. Measurement of RV free wall length may indirectly assess the radial free wall displacement that makes a substantial contribution to RV pump function in PAH patients with dilated RV and preserved tricuspid annular systolic excursion.¹¹

Three-dimensional echocardiography allows direct measurement of RV volumes that have been validated against cardiac magnetic resonance imaging.¹² Three-dimensional imaging technology is progressing rapidly and may become increasingly available for assessment of RV function and geometry in clinical practice.^{13,14} Meanwhile, the predictive value of 2D echocardiography-derived RVESRI needs to be confirmed in other PAH centers.

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From the Division of Cardiology, Tulane University School of Medicine, New Orleans, LA.

Correspondence to Thierry H. Le Jemtel, MD, Tulane University School of Medicine, 1430 Tulane Ave, SL-48, New Orleans, LA 70112. E-mail lejemtel@tulane.edu

(*Circ Cardiovasc Imaging*. 2017;10:e006564.

DOI: 10.1161/CIRCIMAGING.117.006564.)

© 2017 American Heart Association, Inc.

Circ Cardiovasc Imaging is available at
<http://circimaging.ahajournals.org>

DOI: 10.1161/CIRCIMAGING.117.006564

Disclosures

None.

References

- Haddad F, Hunt SA, Rosenthal DN, Murphy DJ. Right ventricular function in cardiovascular disease, part I: anatomy, physiology, aging, and functional assessment of the right ventricle. *Circulation*. 2008;117:1436–1448. doi: 10.1161/CIRCULATIONAHA.107.653576.
- Fernández-Golfín C, Zamorano JL. Three-dimensional echocardiography and right ventricular function: the beauty and the beast? *Circ Cardiovasc Imaging*. 2017;10:e006099. doi: 10.1161/CIRCIMAGING.117.006099.
- Kiernan MS, French AL, DeNofrio D, Parmar YJ, Pham DT, Kapur NK, Pandian NG, Patel AR. Preoperative three-dimensional echocardiography to assess risk of right ventricular failure after left ventricular assist device surgery. *J Card Fail*. 2015;21:189–197. doi: 10.1016/j.cardfail.2014.12.009.
- Bellavia D, Iacovoni A, Scardulla C, Moja L, Pilato M, Kushwaha SS, Senni M, Clemenza F, Agnese V, Falletta C, Romano G, Maalouf J, Dandel M. Prediction of right ventricular failure after ventricular assist device implant: systematic review and meta-analysis of observational studies [published online ahead of print March 31, 2017]. *Eur J Heart Fail*. doi:10.1002/ehj.733.
- Vitarelli A, Mangieri E, Terzano C, Gaudio C, Salsano F, Rosato E, Capotosto L, D'Orazio S, Azzano A, Truscetti G, Cocco N, Ashurov R. Three-dimensional echocardiography and 2D-3D speckle-tracking imaging in chronic pulmonary hypertension: diagnostic accuracy in detecting hemodynamic signs of right ventricular (RV) failure. *J Am Heart Assoc*. 2015;4:e001584. doi: 10.1161/JAHA.114.001584.
- Tedford RJ, Mudd JO, Girgis RE, Mathai SC, Zaiman AL, Houston-Harris T, Boyce D, Kelemen BW, Bacher AC, Shah AA, Hummers LK, Wigley FM, Russell SD, Saggat R, Maughan WL, Hassoun PM, Kass DA. Right ventricular dysfunction in systemic sclerosis-associated pulmonary arterial hypertension. *Circ Heart Fail*. 2013;6:953–963. doi: 10.1161/CIRCHEARTFAILURE.112.000008.
- Smith BC, Dobson G, Dawson D, Charalampopoulos A, Grapsa J, Nihoyannopoulos P. Three-dimensional speckle tracking of the right ventricle: toward optimal quantification of right ventricular dysfunction in pulmonary hypertension. *J Am Coll Cardiol*. 2014;64:41–51. doi: 10.1016/j.jacc.2014.01.084.
- Maffessanti F, Muraru D, Esposito R, Gripari P, Ermacora D, Santoro C, Tamborini G, Galderisi M, Pepi M, Badano LP. Age-, body size-, and sex-specific reference values for right ventricular volumes and ejection fraction by three-dimensional echocardiography: a multicenter echocardiographic study in 507 healthy volunteers. *Circ Cardiovasc Imaging*. 2013;6:700–10. doi: 10.1161/CIRCIMAGING.113.000706.
- Amsallem M, Sweatt AJ, Aymami MC, Kuznetsova T, Selej M, Lu H, Mercier O, Fadel E, Schnittger I, McConnell MV, Rabinovitch M, Zamanian RT, Haddad F. Right heart end-systolic remodeling index strongly predicts outcomes in pulmonary arterial hypertension: comparison with validated models. *Circ Cardiovasc Imaging*. 2017;10:e005771. doi: 10.1161/CIRCIMAGING.116.005771.
- Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W, Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging*. 2015;16:233–270. doi: 10.1093/ehjci/jev014.
- Lakatos B, Tóser Z, Tokodi M, Doronina A, Kosztin A, Muraru D, Badano LP, Kovács A, Merkely B. Quantification of the relative contribution of the different right ventricular wall motion components to right ventricular ejection fraction: the ReVISION method. *Cardiovasc Ultrasound*. 2017;15:8. doi: 10.1186/s12947-017-0100-0.
- Nagata Y, Wu VC, Kado Y, Otani K, Lin FC, Otsuji Y, Negishi K, Takeuchi M. Prognostic value of right ventricular ejection fraction assessed by transthoracic 3D echocardiography. *Circ Cardiovasc Imaging*. 2017;10:e005384. doi: 10.1161/CIRCIMAGING.116.005384.
- Medvedofsky D, Addetia K, Patel AR, Sedlmeier A, Baumann R, Mor-Avi V, Lang RM. Novel approach to three-dimensional echocardiographic quantification of right ventricular volumes and function from focused views. *J Am Soc Echocardiogr*. 2015;28:1222–1231. doi: 10.1016/j.echo.2015.06.013.
- Knight DS, Grasso AE, Quail MA, Muthurangu V, Taylor AM, Toumpanakis C, Caplin ME, Coghlan JG, Davar J. Accuracy and reproducibility of right ventricular quantification in patients with pressure and volume overload using single-beat three-dimensional echocardiography. *J Am Soc Echocardiogr*. 2015;28:363–374. doi: 10.1016/j.echo.2014.10.012.

KEY WORDS: Editorials ■ echocardiography ■ phenotype ■ pulmonary hypertension ■ right ventricle

Respite for 2-Dimensional Right Ventricular Imaging? Rohan Samson and Thierry H. Le Jemtel

Circ Cardiovasc Imaging. 2017;10:

doi: 10.1161/CIRCIMAGING.117.006564

Circulation: Cardiovascular Imaging is published by the American Heart Association, 7272 Greenville Avenue,
Dallas, TX 75231

Copyright © 2017 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-9651. Online ISSN: 1942-0080

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circimaging.ahajournals.org/content/10/6/e006564>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Imaging* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Cardiovascular Imaging* is online at:
<http://circimaging.ahajournals.org/subscriptions/>